

CLOSED IDEALS WITH BOUNDED Δ -WEAK APPROXIMATE IDENTITIES IN SOME CERTAIN BANACH ALGEBRAS

J. LAALI AND M. FOZOUNI

ABSTRACT. It is shown that a locally compact group G is amenable if and only if some certain closed ideals of the Figà-Talamanca-Herz algebra $A_p(G)$ admit bounded Δ -weak approximate identities. Also, similar results are obtained for the function algebras $\mathfrak{L}A(G)$ and $C_0^w(G)$.

1. INTRODUCTION AND PRELIMINARIES

Let A be a Banach algebra, $\Delta(A)$ be the character space of A , that is, the space of all non-zero homomorphisms from A into $\mathbb{C}$ and A^* be the dual space of A consisting of all bounded linear functionals on A . Throughout the paper, A is a commutative and semi-simple Banach algebra, hence $\Delta(A)$ is non-empty.

Let $\{e_\alpha\}$ be a net in Banach algebra A . The net $\{e_\alpha\}$ is called,

- (1) an *approximate identity* if for each $a \in A$, $\|ae_\alpha - a\| \rightarrow 0$,
- (2) a *weak approximate identity* if for each $a \in A$, $|f(ae_\alpha) - f(a)| \rightarrow 0$ for all $f \in A^*$,
- (3) a *Δ -weak approximate identity* if for each $\phi \in \Delta(A)$, $|\phi(e_\alpha) - 1| \rightarrow 0$.

Definition 1. Let A be a Banach algebra. A bounded Δ -weak approximate identity for subspace $E \subseteq A$ is a bounded net $\{e_\alpha\}$ in E such that for each $a \in E$,

$$\lim_{\alpha} |\phi(ae_\alpha) - \phi(a)| = 0 \quad (\phi \in \Delta(A)).$$

For simplicity of notation, let b. Δ -w.a.i stand for a bounded Δ -weak approximate identity.

It was proved that every Banach algebra A with a bounded Δ -weak approximate identity has a bounded approximate identity (b.a.i) and conversely [4, Proposition 33.2].

1991 *Mathematics Subject Classification.* 43A07, 43A15.

Key words and phrases. Character space, amenable group, Figà-Talamanca-Herz algebra, Lebesgue-Fourier algebra.

The notion of a Δ -weak approximate identity introduced and studied in [15] where an example of a Banach algebra which has a Δ -weak approximate identity but does not have an approximate identity, was given. Indeed, if $S = \mathbb{Q}^+$ is the semigroup of positive rationals under addition, it was shown that the semigroup algebra $l^1(S)$ has a b. Δ -w.a.i, but it does not have any bounded or unbounded approximate identity.

As the second example to see the difference between bounded approximate and bounded Δ -weak approximate identities, let $\mathbb{R}$ be the additive real line group and $1 < p \leq \infty$. Put $S_p(\mathbb{R}) = L^1(\mathbb{R}) \cap L^p(\mathbb{R})$ and define the following norm;

$$\|f\|_{S_p} = \max\{\|f\|_1, \|f\|_p\} \quad (f \in S_p(\mathbb{R})).$$

Using [14, Theorem 2.1], we can see that $S_p(\mathbb{R})$ has a b. Δ -w.a.i, but it has no b.a.i. Because we know that $S_p(\mathbb{R})$ is a Segal algebra and it is well-known that a Segal algebra S in $L^1(\mathbb{R})$ has a b.a.i if and only if $S = L^1(\mathbb{R})$. But it is clear that $S_p(\mathbb{R}) \neq L^1(\mathbb{R})$.

These type of approximate identities have some interesting applications, for example; see [9, 16, 22]. In the past decades, B. E. Forrest studied the relations between the amenability of a group G and closed ideals of $A(G)$ and $A_p(G)$ with a b.a.i; see [5–8], and the relations between some properties of G and closed ideals of $A(G)$ with a b. Δ -w.a.i; see [9].

In this paper, we try to improve some of the theorems in [5, 6, 8, 10, 16] with changing b.a.i by b. Δ -w.a.i. As an application, we give the converse of [8, Corollary 4.2] due to B. Forrest, E. Kaniuth, A. T. Lau and N. Spronk.

2. MAIN RESULTS

Let G be a locally compact group. For $1 < p < \infty$, let $A_p(G)$ denote the subspace of $C_0(G)$ consisting of all functions of the form $u = \sum_{i=1}^{\infty} f_i * \tilde{g}_i$ where $f_i \in L^p(G)$, $g_i \in L^q(G)$, $1/p + 1/q = 1$, $\sum_{i=1}^{\infty} \|f_i\|_p \|g_i\|_q < \infty$ and $\tilde{g}(x) = \overline{g(x^{-1})}$ for all $x \in G$. With the pointwise operation and the following norm,

$$\|u\|_{A_p(G)} = \inf \left\{ \sum_{i=1}^{\infty} \|f_i\|_p \|g_i\|_q : u = \sum_{i=1}^{\infty} f_i * \tilde{g}_i \right\},$$

$A_p(G)$ is a Banach algebra called the Figà-Talamanca-Herz algebra. It is clear that $\|u\| \leq \|u\|_{A_p(G)}$ where $\|u\|$ is the uniform norm of $u \in C_0(G)$. By [12, Theorem 3], we know that

$$\Delta(A_p(G)) = \{\phi_x : x \in G\} = G,$$

where ϕ_x is defined by $\phi_x(f) = f(x)$ for each $f \in A_p(G)$.

The dual of the Banach algebra $A_p(G)$ is the Banach space $PM_p(G)$ consisting of all limits of convolution operators associated to bounded measures.

Indeed, $PM_p(G)$ is the w^* -closure of $\lambda_p(L^1(G))$ in $B(L^p(G))$ where λ_p is the left regular representation of G on $L^p(G)$; see [3] for more details.

The group G is said to be *amenable* if, there exists an $m \in L^\infty(G)^*$ such that $m \geq 0$, $m(1) = 1$ and $m(L_x f) = m(f)$ for each $x \in G$ and $f \in L^\infty(G)$ where $L_x f(y) = f(x^{-1}y)$.

Theorem 1. (*Leptin-Herz*) *Let G be a locally compact group and $1 < p < \infty$. Then $A_p(G)$ has a b.a.i if and only if G is amenable*

The proof of the above theorem in the case $p = 2$ is due to Leptin [19] and in general is due to Herz [12].

Forrest and Skantharajah in [9] showed that if G is a discrete group, then $A_2(G) = A(G)$ has a b. Δ -w.a.i if and only if G is amenable. Kaniuth and Ülger in [18, Theorem 5.1], for the first time in our knowledge, announced that $A(G)$ has a b. Δ -w.a.i if and only if G is an amenable group, but the same result holds for the Figà-Talamanca-Herz algebras as follows. The proof is similar to the Fourier algebra case, but we give it for the convenience of reader.

Theorem 2. *Let G be a locally compact group and $1 < p < \infty$. Then $A_p(G)$ has a b. Δ -w.a.i if and only if G is amenable.*

Proof. Let $\{e_\alpha\}$ be a b. Δ -w.a.i for $A_p(G)$ and $e \in A_p(G)^{**}$ be a w^* -cluster point of $\{e_\alpha\}$. So, for each $\phi \in \Delta(A_p(G)) = G$, we have

$$e(\phi) = \lim_{\alpha} \phi(e_\alpha) = 1,$$

because $\{e_\alpha\}$ is a b. Δ -w.a.i for $A_p(G)$. Therefore, by [23, Proposition 2.8] G is weakly closed in $PM_p(G) = A_p(G)^*$. Now, by [2, Corollary 2.8] we conclude that G is an amenable group. $\square$

The following corollary immediately follows from the Leptin-Herz Theorem and Theorem 2.

Corollary 1. *Let G be a locally compact group and $1 < p < \infty$. Then $A_p(G)$ has a b. Δ -w.a.i. if and only if it has a b.a.i.*

The following theorem is a key tool in the sequel.

Theorem 3. *Let A be a Banach algebra, I be a closed two-sided ideal of A which has a b. Δ -w.a.i and the quotient Banach algebra A/I has a b.l.a.i. Then A has a b. Δ -w.a.i.*

Proof. Let $\{e_\alpha\}$ be a b. Δ -w.a.i for I and $\{f_\delta + I\}$ be a b.l.a.i for A/I . We can assume that $\{f_\delta\}$ is bounded. Indeed, since $\{f_\delta + I\}$ is bounded, there exists a positive integer K with $\|f_\delta + I\| < K$ for each δ . So, there exists $y_\delta \in I$ such

that $\|f_\delta + I\| < \|f_\delta + y_\delta\| < K$. Put $f'_\delta = f_\delta + y_\delta$. Clearly, $\{f'_\delta + I\}$ is a b.l.a.i for A/I which $\{f'_\delta\}$ is bounded.

Now, consider the bounded net $\{e_\alpha + f_\delta - e_\alpha f_\delta\}_{(\alpha, \delta)}$. For each $\phi \in \Delta(A)$ we have

$$\phi(e_\alpha + f_\delta - e_\alpha f_\delta) = \phi(e_\alpha) + \phi(f_\delta)(1 - \phi(e_\alpha)) \xrightarrow{(\alpha, \delta)} 1.$$

Therefore, A has a b. Δ -w.a.i. $\square$

Let G be a locally compact group, E be a closed non-empty subset of G and $1 < p < \infty$. Define

$$I_p(E) = \{u \in A_p(G) : u(x) = 0 \text{ for all } x \in E\}.$$

The following result improves [5, Theorem 3.9].

Theorem 4. *Let G be a locally compact group. Then the following assertions are equivalent.*

- (1) G is an amenable group.
- (2) $\ker(\phi)$ has a b. Δ -w.a.i for each $\phi \in \Delta(A_p(G))$.
- (3) $I_p(H)$ has a b. Δ -w.a.i for some closed amenable subgroup H of G .

Proof. (1) $\Rightarrow$ (2): Let G be an amenable group. Then $A_p(G)$ has a b.a.i by Leptin-Herz's Theorem. Now, the result follows from [17, Corollary 2.3].

(2) $\Rightarrow$ (3): Just take $H = \{e\}$, because we know that $I_p(\{e\}) = \ker(\phi_e)$.

(3) $\Rightarrow$ (1): Suppose that $I_p(H)$ for a closed amenable subgroup H of G has a b. Δ -w.a.i. By [21, Lemma 3.19] we know that $A_p(H)$ is isometrically isomorphic to $A_p(G)/I_p(H)$. But $A_p(H)$ has a b.a.i, since H is an amenable group. Therefore, $A_p(G)/I_p(H)$ also has a b.a.i. Now, the result follows from Theorems 3 and 2. $\square$

Forrest in [6, Lemma 3.14] improved [5, Theorem 3.9]. Indeed he showed that G is an amenable group if for some closed proper subgroup H of G , $I_2(H)$ has a b.a.i. Also, using the operator space structure of $A(G)$, it was shown in [8, Theorem 1.5] that G is an amenable group only if $I_2(H)$ for some closed subgroup H of G has a b.a.i.

Now, we give the following result which improves [8, Corollary 1.6] and Theorem 4. Our proof is a mimic of [6, Lemma 3.14].

Theorem 5. *Let G be a locally compact group and $1 < p < \infty$. Then the following are equivalent.*

- (1) G is an amenable group.
- (2) $I_p(H)$ has a b. Δ -w.a.i for some proper closed subgroup H of G .

Proof. In view of [8, Corollary 4.2], only (2) $\Rightarrow$ (1) needs proof.

Let H be a proper closed subgroup of G such that $I_p(H)$ has a b. Δ -w.a.i. We will show that H is an amenable group and this completes the proof by Theorem 4.

Since H is a proper subgroup, there exists $x \in G \setminus H$. On the other hand, the mapping $I_p(H) \rightarrow I_p(xH)$ defined by $u \rightarrow L_x u$ is an isometric isomorphism, because for each $t \in G$ and $f \in A_p(G)$, we have,

$$L_t f \in A_p(G), \quad \|L_t f\|_{A_p(G)} = \|f\|_{A_p(G)}.$$

Therefore, $I_p(xH)$ has a b. Δ -w.a.i which we denote it by (u_α) . For each α , let v_α be the restriction of u_α to H . Using [12, Theorem 1a], we conclude that (v_α) is a bounded net in $A_p(H)$.

Let $\nu \in A_p(H) \cap C_c(H)$ and $K = \text{supp } \nu \subseteq H$. Then there exists a neighborhood V of K in G such that $V \cap xH = \emptyset$, because $K \cap xH = \emptyset$ (otherwise we conclude that x is in H) and G is completely regular by [13, Theorem 8.4] and hence it is a regular topological space. Indeed, for each $y \in xH$, let V_y be a neighborhood of K such that $y \notin V_y$. So, $V = \bigcap_{y \in xH} V_y$ satisfies $V \cap xH = \emptyset$.

By [3, Proposition 1, pp.34] there is $u \in A_p(G)$ such that $u(x) = 1$ for each $x \in K$ and $\text{supp } u \subseteq V$, and by [12, Theorem 1b], there exists a $v \in A_p(G)$ such that $v|_H = \nu$. Now, put $w = vu$. Since $V \cap xH = \emptyset$ and $\text{supp } u \subseteq V$, we have $w \in I_p(xH)$, and since $u(x) = 1$ for each $x \in K$ and $K = \text{supp } \nu$, we have $w|_H = \nu$.

Now, for each $x \in H$, we have

$$\begin{aligned} \lim_{\alpha} |\phi_x(v_\alpha \nu) - \phi_x(\nu)| &= \lim_{\alpha} |v_\alpha(x)\nu(x) - \nu(x)| \\ &= \lim_{\alpha} |u_\alpha(x)w(x) - w(x)| = 0. \end{aligned}$$

Therefore, (v_α) is a b. Δ -w.a.i for $A_p(H)$, since by [3, Corollary 7, pp. 38], $A_p(H) \cap C_c(H)$ is dense in $A_p(H)$. Hence, by Theorem 2, H is amenable. $\square$

As an application of the above theorem, we give the following corollary which is the converse of [8, Corollary 4.2].

Corollary 2. *Let G be a locally compact group, $1 < p < \infty$ and H be a proper closed subgroup of G . If $I_p(H)$ has a b.a.i, then G is amenable.*

Ghahramani and Lau in [10] introduced and studied a new closed ideal of $A(G)$. Indeed, let G be a locally compact group and put

$$\mathfrak{L}A(G) = L^1(G) \cap A(G)$$

with the norm

$$|||f||| = \|f\|_1 + \|f\|_{A(G)}.$$

Clearly $\mathfrak{L}A(G)$ with pointwise multiplication is a commutative Banach algebra with $\Delta(\mathfrak{L}A(G)) = G$ and it is called the *Lebesgue-Fourier algebra* of G .

It was shown that $\mathfrak{L}A(G)$ has a b.a.i if and only if G is a compact group [10, Proposition 2.6]. Now, we give the following result concerning the b. Δ -w.a. identities of this Banach algebra.

Theorem 6. *Let G be a locally compact group.*

- (1) *for each $x \in G$, $\ker(\phi_x) \subseteq \mathfrak{L}A(G)$ has a b. Δ -w.a.i.*
- (2) *$\mathfrak{L}A(G)$ has a b. Δ -w.a.i.*
- (3) *G is amenable.*
- (4) *G is compact.*

Then (1) $\Rightarrow$ (2) $\Rightarrow$ (3) and (4) $\Rightarrow$ (1) hold.

Proof. (1) $\Rightarrow$ (2) : Follows from Theorem 3.

(2) $\Rightarrow$ (3) : Let (u_α) be a b. Δ -w.a.i for $\mathfrak{L}A(G)$, Then (u_α) is a bounded net in $A(G)$. Now, the result follows from Theorem 2 and this fact that $\Delta(\mathfrak{L}A(G)) = G = \Delta(A(G))$.

(4) $\Rightarrow$ (1) : Let G be a compact group. Then by [10, Proposition 2.6], we know that $\mathfrak{L}A(G) = A(G)$. On the other hand, by [17, Example 2.6] for each $x \in G$, $A(G)$ is ϕ_x -amenable. Therefore, the result follows from [17, Proposition 2.2]. $\square$

We do not know whether the implication (3) $\Rightarrow$ (2) in Theorem 6 remains true?

Remark 1. In [11], Granirer gave a (p, r) -version of the Lebesgue-Fourier algebra. Indeed, let $1 < p < \infty$, $1 \leq r \leq \infty$ and put $A_p^r(G) = A_p(G) \cap L^r(G)$ with the norm,

$$\|u\|_{A_p^r(G)} = \|u\|_r + \|u\|_{A_p(G)}.$$

It was shown that $A_p^r(G)$ with pointwise multiplication is a commutative semi-simple Banach algebra such that $\Delta(A_p^r(G)) = G$ and for all $1 \leq r \leq \infty$, $A_p^r(G) = A_p(G)$ if G is a compact group; see [11, Theorem 1, Theorem 2].

Therefore, in view of Theorems 2 and 3, Theorem 6 remains true if we replace $\mathfrak{L}A(G)$ with $A_p^r(G)$.

Remark 2. Runde in [20], by using the canonical operator space structure of $L^p(G)$, introduced and studied the algebra $OA_p(G)$ for $1 < p < \infty$, the *operator Figà-Talamanca-Herz algebra*. It was shown that $A_p(G) \subseteq OA_p(G)$ [20, Remark 4. pp. 159] and $OA_p(G)$ has a b.a.i if and only if G is an amenable group [20, Theorem 4.10]. That would be an interesting question: Are the preceding results remain true if we replace $A_p(G)$ by $OA_p(G)$?

Now, let G be a locally compact group and $w : G \rightarrow \mathbb{R}$ be an upper semi-continuous function such that $w(x) \geq 1$ for each $x \in G$. Put

$$C_0^w(G) = \{f \in C(G) : fw \in C_0(G)\}.$$

It is clear that $C_0^w(G)$ with pointwise operation and weighted supremum norm defined by

$$\|f\|_w = \sup_{x \in G} |f(x)|w(x) \quad (f \in C_0^w(G)),$$

is a commutative Banach algebra such that $\Delta(C_0^w(G)) = \Delta(C_0(G)) = G$; see [16, Section 4.3].

Theorem 7. *Let G be a locally compact group and $t \in G$. Then the following are equivalent.*

- (1) $\ker(\phi_t)$ has a b.a.i.
- (2) $\ker(\phi_t)$ has a b. Δ -w.a.i.
- (3) $C_0^w(G)$ has a b. Δ -w.a.i.
- (4) w is bounded.

Proof. (1) $\Rightarrow$ (2) : This part is clear. Applying Theorem 3, we conclude (2) $\Rightarrow$ (3). In view of [16, Corollary 4.7], we have (3) $\Rightarrow$ (4). Therefore, we only show (4) $\Rightarrow$ (1).

Suppose that w is bounded by $M > 0$. Let $\mathcal{V} = \{V_\alpha\}_{\alpha \in \Gamma}$ be a neighborhood base at t directed by the reverse inclusion. For each $\alpha \in \Gamma$, by the Urysohn Lemma, there exists a continuous function $f_\alpha : X \rightarrow [0, 1]$ such that $f_\alpha(t) = 1$ and $\text{supp} f_\alpha \subseteq V_\alpha$.

Let $\epsilon > 0$ and $g \in C_0^w(G)$. For $\epsilon' = \epsilon/2M$, there exists a neighborhood V of t such that,

$$|g(y) - g(t)| < \epsilon' \quad (y \in V).$$

On the other hand, let $\alpha_0 \in \Gamma$ be such that $V_{\alpha_0} \subseteq V$. Therefore, there exists $x_0 \in G$ such that

$$\begin{aligned} \|gf_\alpha - \phi_t(g)f_\alpha\|_w &= \sup_{x \in G} |g(x)f_{\alpha_0}(x) - g(t)f_{\alpha_0}(x)|w(x) \\ &< |g(x_0)f_{\alpha_0}(x_0) - g(t)f_{\alpha_0}(x_0)|w(x_0) + \epsilon/2 \\ &< M\epsilon' + \epsilon/2 = \epsilon. \end{aligned}$$

Therefore, for each $g \in C_0^w(G)$, $\|gf_\alpha - g(t)f_\alpha\|_w \rightarrow 0$. Hence, by [17, Theorem 1.4, Proposition 2.2] and [1, Corollary 3.6] we conclude that $\ker(\phi_t)$ has a b.a.i. $\square$

ACKNOWLEDGEMENT

The authors are grateful to the referee of the paper for his (her) invaluable comments which improved the presentation of the paper, especially for giving a shorter proof for Theorem 3.

REFERENCES

- [1] ARHIPPAINEN, J. AND KAUPPI, J.: *Generalization of the B^* -algebra $(C_0(x), \|\cdot\|_\infty)$* , Math. Nachr., **282** (2009), No. 1, 7–15.
- [2] CHOU, C. AND XU, G.: *The weak closure of the set of left translation operators*, Proc. Amer. Math. Soc., **127** (1999), 465–470.
- [3] DERIGHETTI, A.: *Convolution Operators on Groups*, Springer-Verlag, 2011.
- [4] DORAN, R. S. AND WICHMANN, J.: *Approximate Identities and Factorization in Banach Modules*, vol. 768, Springer-Verlag, 1979.
- [5] FORREST, B. E.: *Amenability and bounded approximate identities in ideals of $A(G)$* , Illinois J. Math., **34** (1990), 1–25.
- [6] FORREST, B. E.: *Amenability and ideals in $A(G)$* , J. Austral. Math. Soc. (Series A), **53** (1992), 143–155.
- [7] FORREST, B. E.: *Amenability and the structure of the algebras $A_p(G)$* , Trans. Amer. Math. Soc., **343** (1994), No. 1, 233–243.
- [8] FORREST, B. E., KANIUTH, E., LAU, A. T.-M., AND SPRONK, N.: *Ideals with bounded approximate identities in Fourier algebras*, J. Funct. Anal., **203** (2003), 286–304.
- [9] FORREST, B. E. AND SKANTHARAJAH, M.: *A note on a type of approximate identity in the Fourier algebra*, Proc. Amer. Math. Soc., **120** (1994), No. 2, 651–652.
- [10] GHAHRAMANI, F. AND LAU, A. T.-M.: *Weak amenability of certain classes of Banach algebras without bounded approximate identities*, Math. Proc. Camb. Phil. Soc., **133** (2002), 357–371.
- [11] GRANIRER, E. E.: *The Figà-Talamanca-Herz-Lebesgue Banach algebras $A_p^r(G) = A_p(G) \cap L^r(G)$* , Math. Proc. Camb. Phil. Soc., **140** (2006), 401–416.
- [12] HERZ, C.: *Harmonic synthesis for subgroups*, Ann. Inst. Fourier (Grenoble), **XXIII** (1973), No. 3, 91–123.
- [13] HEWITT, E. AND ROSS, K.: *Abstract Harmonic Analysis I*, Springer-Verlag, Berlin, 1979, 2nd edn.
- [14] INOUE, J. AND TAKAHASI, S.-E.: *Constructions of bounded weak approximate identities for Segal algebras on LCA groups*, Acta Sci. Math. (Szeged), **66** (2000), No. 1-2, 257–271.
- [15] JONES, C. A. AND LAHR, C. D.: *Weak and norm approximate identities are different*, Pacific J. Math., **72** (1977), No. 1, 99–104.
- [16] KAMALI, Z. AND BAM, M. L.: *Bochner-Schoenberg-Eberlein property for abstract Segal algebras*, Proc. Japan Acad. Ser. A., **89** (2013), 107–110.
- [17] KANIUTH, A. T.-M., LAU, E. AND PYM, J.: *On ϕ -amenability of Banach algebras*, Math. Proc. Camb. Phil. Soc., **144** (2008), 85–96.
- [18] KANIUTH, E. AND ÜLGER, A.: *The Bochner-Schoenberg-Eberlian property for commutative Banach algebras, especially Fourier-Stieltjes algebras*, Trans. Amer. Math. Soc., **362** (2010), 4331–4356.
- [19] LEPTIN, H.: *Sur l’algebre de Fourier d’un groupe localement compact*, R. Acad. Sci. Paris, Ser. A., **266** (1968), 1180–1182.
- [20] RUNDE, V.: *Operator Figà-Talamanca-Herz algebras*, Studia Math., **155** (2003), No. 2, 153–170.
- [21] SANGANI MONFARED, M.: *Extensions and isomorphisms for the generalized Fourier algebras of a locally compact group*, J. Funct. Anal., **198** (2003), 413–444.
- [22] TAKAHASI, S.-E. AND HATORI, O.: *Commutative Banach algebras which satisfy a Bochner-Schoenberg-Eberlein type-theorem*, Proc. Amer. Math. Soc., **110** (1990), No. 1, 149–158.

- [23] ÜLGER, A.: *Some results about the spectrum of commutative Banach algebras under the weak topology and applications*, Monatsh. Math., **121** (1996), 353–379.

Javad Laali, KHARAZMI UNIVERSITY, FACULTY OF MATHEMATICAL AND COMPUTER SCIENCE, DEPARTMENT OF MATHEMATICS, 50 TALEGHANI AVENUE, 15618, TEHRAN, IRAN
E-mail address: laali@khu.ac.ir

Mohammad Fozouni, GONBAD KAVOUS UNIVERSITY, FACULTY OF SCIENCES AND ENGINEERING, DEPARTMENT OF MATHEMATICS, GONBAD KAVOUS, IRAN
E-mail address: fozouni@hotmail.com